

AD-A087 208

NAVAL RESEARCH LAB WASHINGTON DC

F/G 4/1

QUASILINEAR SCATTERING FROM WAVES DRIVEN BY BEAM-PLASMA INSTABILITY--ETC(U)

APR 80 S H BRECHT, P J PALMADESSO

UNCLASSIFIED

NRL-MR-4203

SBIF-AD-E000 474

NL

[DE]
ADA
D-47408



END
DATE
FILMED
9-80
DTIC

ADA 087208

Quantum Electrodynamics of Photon-Photon Interactions

S. H. HEDGECOCK

Physics Department, Los
Alamos National Lab.


AND

P. J. RAYNAUD

Department of Physics, University of
California, Santa Barbara

April 21, 1980

This report was sponsored by the Defense Nuclear Agency under contract
Work Unit 44, Title "Gamma Gamma"



DNAC

COPI

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER NRL Memorandum Report 4203	2. GOVT ACCESSION NO. AD-A087206	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) QUASILINEAR SCATTERING FROM WAVES DRIVEN BY BEAM-PLASMA INSTABILITIES		5. TYPE OF REPORT & PERIOD COVERED Interim report on a continuing NRL problem.
7. AUTHOR(s) S.H. Brecht* and P.J. Palmadesso		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Naval Research Laboratory Washington, DC 20375		8. CONTRACT OR GRANT NUMBER(s)
11. CONTROLLING OFFICE NAME AND ADDRESS Defense Nuclear Agency Washington, DC 20305		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 67-0914-0-0 S99QAXHC062/44
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE April 21, 1980
		13. NUMBER OF PAGES 34
		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES *Present address: Science Applications, Inc., McLean, Virginia 22102. This research was sponsored by the Defense Nuclear Agency under subtask S99QAXHC062, work unit code 44, and work unit title Starfish Optical.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Starfish southern conjugate data Quasilinear effects Ion cyclotron instabilities Beam-plasma interaction		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The Starfish southern conjugate rebrightening is explained by velocity diffusion perpendicular to the earth's magnetic field line. The diffusion is caused by particle scattering off of the harmonics of ion cy- clotron waves commonly called Bernstein modes. Quasilinear estimates of the wave fluctuation levels indicate that the debris streaming down the field lines will excite these waves and then be scattered by them.		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-014-6601

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

CONTENTS

INTRODUCTION	1
LINEAR CALCULATION	4
QUASILINEAR CALCULATION	9
DISCUSSION	15
CONCLUSIONS	18
REFERENCES	20
DISTRIBUTION LIST	27

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DDC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/ _____	
Availability Codes	
Dist.	Avail and/or special
A	

QUASILINEAR SCATTERING FROM WAVES DRIVEN BY
BEAM-PLASMA INSTABILITIES

INTRODUCTION

Test data taken during the Starfish event of 1962 displays a marked second brightening of the atmosphere in the southern conjugate region. This brightening followed the initial luminescence in the southern conjugate region caused by the deposition of the initial debris by approximately 10 seconds.

The explanation that appears most probable involves scattering a fraction of the streaming debris out of the loss cone as the debris proceeds toward the southern conjugate region. The scattering sources are waves excited to high levels driven by the initial fast debris. The scattered debris would then mirror in the southern conjugate and stream toward the magnetic bubble. Some fraction would be scattered back into the loss cone region by the loss cone instability. The remainder would mirror off of the bubble and return toward the southern conjugate region. During the return the loss cone region could be filled by loss cone instabilities and a second precipitation of debris particles could occur. The second precipitation would involve a greater proportion of the distribution because of nonlinear flattening of the distribution function.

In this report the possibility of exciting the waves which are predicted to perform the scattering is addressed. It will be shown by use of a numerical code and quasilinear estimates that indeed waves can be excited and will grow to sufficient levels to cause quasilinear diffusion of the streaming debris distribution function.

The code employed is a linear code which has been used to study beam-plasma instabilities in tokamaks. The circumstances of the problem are similar to a beam-plasma interaction, the general type of which, have been studied for a number of years in the fusion community.¹⁻⁵ The linear code calculates the energy transfer between the beam and waves of interest and the waves with the background plasma to determine the growth rates and energy transferred to the wave. From these calculations quasilinear estimates are made to determine the amount of scattering taking place.

There are a large number of waves which may be excited by a beam of high energy ions.⁴ Among them are the ion cyclotron wave, ion Bernstein wave, ion acoustic wave, low-hybrid waves and shear Alfvén waves. Several of these waves are of high frequency and candidates for study. However, the ion cyclotron wave and its higher harmonics, usually called the ion Bernstein wave, are chosen for study because these waves have been predicted theoretically¹ and shown experimentally⁶⁻¹⁰ to be excited during beam plasma interactions. The code employed for this work was successful in predicting quite accurately these instabilities in a variety of plasma devices from Q machines, to Tokamaks such as the French tokamak, TFR,¹⁰ and PLT, the Princeton Large Torus.

In the following section the details for calculating the linear growth rates for the waves will be briefly discussed. The quasilinear estimates are presented in the following section and finally the conclusion of these calculations are presented.

LINEAR CALCULATION

The linear calculation is based on a perturbation method where the real frequency is assumed larger than the imaginary, $\omega_r > \omega_i$.^{4,11} The code calculates the rate of change of the wave energy by computing a time averaged $\underline{J} \cdot \underline{E}$ for each species given by the following formula

$$\frac{dW_k}{dt} = \langle \underline{J} \cdot \underline{E} \rangle \equiv \sum_j L_j \equiv \sum_j n_j \mathcal{L}_j \quad (1)$$

where the sum j is over the plasma species. Physically the process of energy transfer takes place through resonant processes either Landau cyclotron or anomalous cyclotron resonances. The beam transfers energy to the wave and background plasma. The wave in turn transfers energy to the background electron or ions. If the beam pumping rate is larger than the damping rates the wave grows.

Typically, the instability is parameterized by the amount of beam density necessary to drive the wave unstable, n_b . From Eq. (1) a marginal stability criterion can be established. For marginal stability

$$n_i \mathcal{L}_i + n_e \mathcal{L}_e + n_b \mathcal{L}_b = 0 \quad (2)$$

where n_j are the densities of the species and \mathcal{L}_j is the power transferred per particle. One then obtains

$$\left(\frac{n_b}{n_e} \right)_{\text{critical}} = - \frac{\mathcal{L}_e + n_i \mathcal{L}_i / n_e}{\mathcal{L}_b} \quad (3)$$

and

$$\gamma = \frac{1}{2W_k} \frac{dW_k}{dt} = \frac{n_e \mathcal{L}_b}{W_k} \left(\frac{n_b}{n_e} - \left(\frac{n_b}{n_e} \right)_{\text{crit.}} \right) \quad (4)$$

where Eq. (4) is the equation for the growth rate. If n_b is known then the code calculates the power transfer per particle for the beam, \mathcal{L}_b , and the wave energy for the wave being studied to produce the growth rate. Obviously if $n_b/n_e < (n_b/n_e)_{\text{crit.}}$ the mode is damped. The power transfer function is calculated from the following formulas;^{1,4}

$$\mathcal{L}_j = \frac{-e^2 z_j}{4 |k_{||}| v_j^3 m_j} \sum_{\ell=-\infty}^{\infty} |M_j^\ell| \epsilon_j^\ell \quad (5)$$

where

$$M_j^\ell = \frac{\ell \omega_{cj} E_{\perp}}{k_{\perp}} + \frac{\omega - \ell \omega_{cj}}{k_{||}} E_{||} \quad (6)$$

and

$$\epsilon_j^\ell = 4\pi^2 \int_0^\infty dv_{\perp} v_{\perp} J_\ell^2 \left(\frac{k_{\perp} v_{\perp}}{2\omega_{cj}} \right) \left[\left(1 - \frac{\ell \omega_{cj}}{\omega} \right) \frac{\partial F_j}{\partial v_{||}^2} + \frac{\ell \omega_{cj}}{\omega} \frac{\partial F_j}{\partial v_{\perp}^2} \right] \quad (7)$$

$v_{||} = \frac{c_j^\ell}{\omega}$

with

$$c_j^\ell \equiv (\omega - \ell\omega_{cj})/k_{||}$$

For Maxwellian Species

$$\mathcal{L}_{j=e,i} = \frac{Z_j^2 e^2}{2|k_{||}|v_{jm_j}^3} \sum_{\ell=-\infty}^{\infty} |M_j^\ell|^2 I_\ell(b) \bar{e}^b e^{-(\omega - \ell\omega_{cj})^2/k_{||}^2 v_j^2} \quad (8)$$

where $I(b)$ is the modified Bessel function, $b = \frac{1}{2} \left(\frac{kv}{\omega_{cj}} \right)^2$ and $v_j = \left(2T_j/m_j \right)^{1/2}$.

As can be seen from Eq. (7) in order to perform the calculations one needs to know the distribution function for the beam, F_b . One could assume a monoenergetic beam of particles streaming down the field lines but while such a distribution will produce unstable waves it does not appear realistic.

Noting that the debris streaming down the field lines has already interacted with the coupling shell through a variety of instabilities, i.e., loss cone and mirror instabilities, one needs a more sophisticated model. The model must also consider the effects of velocity dispersion because the excitation of the scattering wave can occur at considerable distances from the burst point. For a simple model of these effects one can use the Fokker-Planck Equation¹ to obtain a distribution which will represent the fact that the debris particles have been scattered in velocity space. Inclusion of a source term in the Fokker-Planck equation

allows one to model the velocity dispersion. That is, particles are introduced into velocity space at a given location. The results of running the Fokker-Planck section of the code is shown in Figure 1. The beam was injected at 200 keV or a velocity of 1.18×10^8 cm/sec. The distribution was allowed to evolve until an appropriate spread in velocity was obtained. As can be seen in Figure 1 the resulting distribution function the particles have a pitch angle ratio of about 10 to 1, $v_{||}$ to v_{\perp} , consistent with the mirror ratio of the magnetic bubble. In addition, the particle density is weighted toward the high velocity side which is the effect produced by velocity dispersion along the magnetic field line. The slow edge of the distribution has a parallel velocity near 7.3×10^7 cm/sec.

This distribution while not having the exact details of the debris distribution function appears to have the quantitative details which are of interest. In the calculations to be discussed, this is the distribution employed. It represents, we feel, a conservative estimate due to the smoothness and the way we introduced the beam. Only one-quarter of the actual beam density was obtained so the actual gradients are in fact smaller than might be expected.

It should be noted that the effect of the velocity dispersion is very important because the instability grows in the positive slope region of the instability. This characteristic has a definite effect in determining where the velocity will be excited and what part of the total debris spectrum will be affected. These effects will be discussed later in this report.

Upon running the linear code it was found that, while the fundamental electrostatic mode of the ion cyclotron wave could be excited, very little energy was transferred into this mode. Therefore growth rates were quite small and insufficient for the anticipated pulse width of the debris. The details of the dispersion relation for the ion cyclotron wave and its higher harmonics, the ion Bernstein waves, can be found in Reference 1.

It was found that the ion Bernstein waves could be excited with quite large growth rates. These rates approached the real frequency of the mode when n_b was taken to be 20% of n_e . The parameters used for these calculations are shown in Table I. Two harmonics of the wave were considered. In the first wave to be studied the second cyclotron harmonic of the beam cyclotron frequency resonates with the wave near the second harmonic of the plasma cyclotron frequency, i.e.,: $\omega_{cb} \approx \omega_{ci}$ where $\ell_b = \ell = 2$. In the case of exciting the third harmonic of the ion Bernstein mode the resonate match was $\ell_b = 2, \ell = 3$.

Both waves were found to be easily excited because the resonant nature of the modes minimizes the Landau and cyclotron damping of the modes. There was considerable energy transfer from the beam to the wave and the growth rates approached ω_{ci} , the limits of the code. Table II show typical results of the second harmonic calculation. It can be seen from Table II that the typical scale for the wavelengths are about 6 km in the perpendicular direction and 30 km in the parallel direction. The group velocities appear to be slow enough so that the excitation process will occur on time scales faster than the energy transport. In summary

the results from the linear calculations indicate that waves of the ion Bernstein type will be excited and so will some of the higher frequency waves.

QUASILINEAR CALCULATION

In this section the quasilinear estimates for the pitch angle scattering of the streaming debris are addressed. The purpose of this discussion is to ascertain whether or not quasilinear effects would be important. It should be noted from the outset that to perform this calculation with complete accuracy one needs to do the problem numerically in two dimensions. However, the approximation to one dimensional diffusion is reasonably good as the perpendicular electron field of the wave, E_{\perp} , is considerably larger than the parallel E_{\parallel} , $E_{\perp} \gg E_{\parallel}$. The general approach for quasilinear calculations can be found in Reference 12.

As mentioned, the waves being considered are electrostatic in nature. This allows one to write the equation for the quasilinear behavior of the beam distribution as

$$\begin{aligned} \frac{\partial f_b}{\partial t} = & \pi \left(\frac{e_b}{m_b} \right)^2 \sum_{k_{\parallel} k_{\perp}} \left(k_{\parallel} \frac{\partial}{\partial v_{\parallel}} + \frac{\ell \omega_{cb}}{v_{\perp}} \frac{\partial}{\partial v_{\perp}} \right) |\phi_k|^2 \delta(\omega - \ell \omega_{cb} - k_{\parallel} v_{\parallel}) \\ & \times J_{\ell}^2(k_{\perp} v_{\perp} / \omega_{cb}) \left(k_{\parallel} \frac{\partial f_b}{\partial v_{\parallel}} + \frac{\ell \omega_{cb}}{v_{\perp}} \frac{\partial f_b}{\partial v_{\perp}} \right), \end{aligned} \quad (9)$$

where J_ℓ is the Bessel function and ϕ_k is the electrostatic potential. For the unstable modes the parallel wave electric field is sufficiently small that the quasilinear diffusion reduces to a diffusion in v_\perp at a constant v_\parallel given by

$$\left. \frac{\partial f_b}{\partial \tau} \right|_q = \frac{1}{v_\perp} \frac{\partial}{\partial v_\perp} \left[v_\perp D(v_\perp, v_\parallel) \right] \frac{\partial f_b}{\partial v_\perp} \sim \frac{f_b}{\tau_q}, \quad (10)$$

where

$$D(v_\perp, v_\parallel) = \frac{n_b^2}{m_b^2} \left(\frac{\ell \omega_{cb}}{v_\perp} \right)^2 \sum_k |\phi_k|^2 J_\ell^2 \left(\frac{k_\perp v_\perp}{\omega_{cb}} \right) \delta(\omega_k - \ell \omega_{cb} - k_\parallel v_\parallel). \quad (11)$$

Equation (10) defines the quasilinear time, i.e., the time scale upon which one expects quasilinear effects to act. The sum over the resonant modes is related to the sum over the fluctuation spectrum through the correlation time given by $\tau_c = 1/\Delta k_\parallel v_b = (\Delta v_b/v_b)/\delta\omega$ where $\Delta v_b/v_b$ is the relative width of the parallel velocity distribution and $\delta\omega = \omega - \ell \omega_{cb}$. This time scale is a measure of the time the beam will stay in resonance with the wave. Although the time τ_c can be long compared to the wave period, one finds that the quasilinear conditions remain valid for the noise levels estimated here for many e-folding times.

One can define the ratio of rf wave energy density to the thermal plasma density by

$$\Gamma_{rf} = \frac{W_k}{\frac{3}{2} (n_e T_e + n_i T_i)} \quad (12)$$

where the wave energy, W_k , is defined in Reference 1 as

$$W_k \sim \frac{2\omega_{pi}^2}{k_{\perp}^2 v_i^2} \left(\frac{\ell\omega_{ci}}{\omega - \ell\omega_{ci}} \right)^2 \frac{I_{\ell}(b) e^{-b}}{16\pi} |E_k|^2, \quad (13)$$

$$\text{where } b \equiv \left(\frac{k_{\perp} v_i}{\omega_{ci}} \right)^2 / 2.$$

For this case $E_k = E_{\perp}$ and it is assumed that $T_e = T_i = T$ for the rest of the calculation. Substituting Eq. (13) into Eq. (12) returns the following expression

$$\Gamma_{rf} = \frac{1}{12} \sum_k \left| \frac{e\phi}{T} \right|^2 \frac{(\ell\omega_{cb})^2}{(\omega - \ell\omega_{cb})^2} I_{\ell}(b) e^{-b}. \quad (14)$$

Using Eqs. (10) and (11) as well as the correlation time leads to the expression for the quasilinear time

$$\frac{1}{\tau_q} \sim \frac{\pi e_b^2 v_{b\parallel}}{m_b^2 \Delta v_{b\parallel} \Delta v_{b\perp}^2 v_{b\perp}^2} \frac{(\ell\omega_{cb})^2}{(\omega - \ell\omega_{cb})^2} J_{\ell} \left(\frac{k_{\perp} v_i}{\omega_{cb}} \right) \sum_{k_{\perp}} |\phi_k|^2. \quad (15)$$

From Eqs. (14) and (15) one obtains the final form for the quasilinear time

$$\frac{1}{\tau_q} = 3\pi\omega_{cb} \Gamma_{rf} \left(\frac{T}{E_b} \right)^2 G_b \frac{\omega - \ell\omega_{cb}}{\omega_{cb}} \frac{J_\ell^2 (k_\perp v_\perp / \omega_{cb})}{I_\ell(b) \bar{e}^b} e^{2\gamma t}, \quad (16)$$

where

$$G_b \equiv v_b^4 v_{b||} / v_{b\perp}^2 \Delta v_{b||} \Delta v_{b\perp}^2$$

measures the localization of the fast ion distribution velocity space. It is assumed that the fluctuation potential begins from thermal levels therefore W_k (thermal) = $(2\pi)^{-3} T \Delta^3 k$ and using Eq. (13) the electric field at the thermal level is estimated to be

$$E_1^2 = \frac{T \Delta^3 k 16\pi(\omega - \ell\omega_{ci})^2}{I_\ell(b) \bar{e}^b \ell^2 \omega_{pi}^2} e^{2\gamma t_0}, \quad t_0 = 0. \quad (17)$$

One last relation must be determined, that is the growth time necessary for the mode to grow to nonlinear or saturated levels. This can be estimated by recalling that trapping effects will occur when the noise level has grown sufficiently. Typically this would occur when the trapping velocity $(e\tilde{\phi}/m_b)^{1/2}$ becomes equal to $\Delta v_{b||}$ or equivalently when the resonant fast ion bounce frequency, $k_{||}(e\tilde{\phi}/m_b)^{1/2}$, becomes equal to the decorrelation rate $1/\tau_c$. Applying this condition to Eq. (14) one arrives at the relation

$$\Gamma_{\text{rf}}(\text{trap}) \sim \frac{1}{12} \left(\frac{\Delta v_{b||}}{c_s} \right)^4 \left(\frac{\ell \omega_{cb}}{\omega - \ell \omega_{cb}} \right)^2 I_\ell(b) e^{-b} \quad (18)$$

with c_s defined as the sound speed. At this point one can estimate the time it will take the mode to encounter nonlinear effect with the relation,

$$\tau_s \sim \frac{1}{2\gamma} \ln \left(\frac{\Gamma_{\text{rf}}(\text{trap})}{\Gamma_{\text{rf}}(\text{thermal})} \right) . \quad (19)$$

Using the values contained in Tables I, II, and III, Eqs. (16) and (19) can be evaluated. The quasilinear time estimate and the trapping time estimate can be compared to one another and with the correlation time to determine if the quasilinear effects have been self-consistently estimated.

Equations (16) and (19) along with the relation for the correlation time were numerically evaluated. The correlation time, τ_c , was found to be in the range of 0.025 to 0.1 sec. This implies that the positive slope of the distribution function such as shown in Figure 1 will be resonant with the wave for times of order 0.1 sec. The quasilinear time, τ_q , which is the measure to determine if strong pitch angle scattering will occur on a time scale comparable with or faster than the correlation time, is found to be on order 10^{-3} sec after the perturbed potential has grown for ~ 0.05 to 0.1 sec. The higher harmonic modes reach the point of rapid quasilinear diffusion in shorter periods. Finally, the saturation time, τ_s , is estimated to be approximately 0.05 to 0.1 sec.

The results of evaluating the relations for τ_c , τ_q and τ_s is that τ_c and τ_s operate on comparable time scales and that on these time scales the wave potential has grown to sufficient levels to produce quite rapid quasilinear effects as reflected by the small values of τ_q at times on the order of τ_c or τ_s .

In Figure 2, the results of evaluating the equation below are shown

$$v_{\perp}(t) = v_{\perp}(0) + \frac{e}{m_b \gamma} E_{\perp 0} (e^{\gamma t} - 1) \quad (20)$$

where $E_{\perp 0}$ is the thermal level of the perpendicular electric field. In this figure plots are made for $\gamma \sim 2\omega_{ci}$ and $\gamma \sim 3\omega_{ci}$. They indicate that $v_{\perp}(t)$ will reach magnitudes large enough to remove the debris from the loss cone region on the time scale of the estimated saturation time, τ_s . From Figure 2, it can be seen that although the third harmonic wave begins at a lower fluctuation level, due to its faster growth rate, it reaches levels of sufficient size to cause the pitch angle scattering more rapidly than the second harmonic. It is expected that waves with faster growth rates would continue this trend.

Both the second and third harmonic waves appear in Figure 2 to produce sufficient pitch angle scattering to explain the second brightening phenomena. The estimates indicate that the time scales for the wave to grow and reach at least saturation amplitudes are faster than the pulse width of debris.

DISCUSSION

Due to the very complicated relations between the wave dispersion relations and the beam distribution function in this calculation and the lack of knowledge about the details of the time evolution of the debris spectrum, it is impossible to make statements concerning the exact region of the debris distribution function which will excite the waves. Estimates can be made, however, which will indicate probable areas of interest.

Before these estimates can be made some details of the wave dispersion relation must be presented. It was found during the numerical calculation that the frequency at any k_{\perp} valid with the parameter regime of the calculation was independent of $k_{||}$. It was also found in this same parameter regime that the frequency was slowly varying in k_{\perp} . Therefore, to good accuracy over the full range of possible $v_{b||}$ one could consider the frequency shift, $\omega - \ell\omega_{ci}$, to be a constant.

During the course of the calculation it was found that the modes that could be excited had a very limited range in $k_{||}$ and that this preferred parallel wavelength was about 30 km. To some extent this particular wavelength is a function of the distribution used but the narrowness of the spectrum is probably independent of it. As can be seen in Figure 1, there was a fair range of $v_{||}$ space in which the distribution had a positive slope but the modes discussed here were the only ones undergoing energy transfer. Using the resonant condition for beam wave energy transfer, $v_{b||} = (\omega - \ell\omega_{cb})/k_{||}$, and noting that the scale height of the atmosphere may limit the length of any given mode, one can estimate an upper bound on the parallel velocity which might

excite waves. With the scale height ~ 100 km and recalling the $\omega - \omega_{cb} \sim \text{constant}$ the maximum parallel beam velocity is estimated to be about 2.3×10^8 cm/sec. At equatorial latitudes the scale height argument does not apply and one may find longer wavelengths thus higher parallel velocities. One can make additional estimates by noting that the mode which appears to be the most unstable in this sample calculation has a parallel wavelength of about 30 km and that the correlation time, where $\tau_c \sim \Delta v_b / (v_b \delta\omega)$, has a range of about 0.025 sec to 0.1 sec. With these parameters the range of velocities which might excite the waves is found to be about 6×10^7 cm/sec to 1.2×10^8 cm/sec. However, once the waves have been excited debris of all velocities would undergo strong pitch angle scattering. Clearly particles with higher velocities than those exciting the wave would also be scattered by these waves but the proportion of particles scattered at any given energy would be lower. This occurs because the high energy particles emitted from the coupling shell at the earliest times would encounter low levels of wave turbulence while high energy particles emitted later would see high fluctuation levels. The exact timing of such occurrences is difficult to ascertain.

The time of arrival for the particles with velocities in the range 6×10^7 to 2.3×10^8 cm/sec is the southern conjugate region (SCR) can be easily estimated. Using 4000 km as a canonical distance, one finds that these particles will arrive in the SCR in the time frame of 1.8 to 7 sec. However, since these particles are expected to excite the waves they represent the sections of the distribution which will be most drastically isotropized by the waves. This would remove as much as 50% of the debris population in this velocity range from the loss cone and would account for the second brightening.

While this discussion has been directed toward specific set of Bernstein modes, one could expect a spectrum of both varying wavelengths and different frequency harmonics to be excited. The higher harmonic modes, $\ell > 2$, would be excited by the faster portions of the streaming debris and will be scattered by these modes. The fluctuation levels may be expected to be less than the modes discussed at length here due to energy coupling difficulties encountered by the higher harmonic modes.¹ In any case the range of the spectrum will be greatly effected by the exact details of the debris distribution as a function of time and space. As a final comment estimates made from the SCORPIO code runs indicate that there would be sufficient beam density to excite these waves in the parameter region discussed.

CONCLUSIONS

The results of the linear calculations indicate that the ion Bernstein modes near the second and third harmonic of the plasma cyclotron frequency will be excited by the streaming debris. Modes of higher frequency may also be excited but were not considered in these calculations.

Quasilinear estimates indicate that quasilinear effects will occur on time scales shorter than the debris pulse width ~ 1 sec. Using simple estimates for the growth rates and the amplitude of the perpendicular electric field as a function of time, it is found that considerable diffusion of the debris distribution function will occur on time scales comparable with the saturation time, i.e., ~ 0.1 sec.

The pitch angle scattering predicted from the quasilinear estimates is more than sufficient to remove a significant fraction of the streaming debris particles from the loss cone. From the estimates made of the range of possible beam velocities that might excite the wave and undergo pitch angle scattering it was found that a large percentage of the total debris distribution was susceptible to these instabilities.

The estimates appear to be in agreement with the data in the SCR. The second brightening in this region appeared to be nearly as bright as the first indicating something approaching equal partition of the energy between the two depositions. From these calculations most of the debris distribution would be susceptible to the instabilities and one then has the possibility of up to 50% of the debris being mirrored. This of course would be the upper bound and a more realistic number probably involves factors of two. An additional phenomena was observed during

the SCR deposition. It appeared on the film that the altitude of deposition between the first and second brightening had changed. The region of luminescence caused by the second deposition appeared at a higher altitude. These results would indicate that the energy of the most energetic particles in the debris being deposited was less than at the time of the first deposition. This again appears to be consistent with the effects of the instability. As the waves are excited, the particles must lose energy. This would have the effect of decreasing the energy of the portions of the distribution which is exciting the waves. Typically the higher energy particles.

In summary, from the linear and quasilinear calculations it is certain that the ion Bernstein waves can be excited. It appears that they can be excited to sufficient wave amplitude to cause strong pitch angle scattering and explain the second brightening seen in the southern conjugate region during the Starfish event. It also seems clear that nonlinear effects may make the phenomena discussed here even more pronounced if flattening of the distributions occurs. This flattening would inhibit the excitation of these waves after the isotropization. The result of this would be that upon subsequent bounces the particles would not be scattered out of the loss cone by these instabilities and would then precipitate for the most part upon return to the SCR, with particles being returned to the loss cone region by loss cone instabilities. Work on a problem similar to this is discussed in References 13 and 14 where the authors have addressed the problem of electron trapping in the radiation belts.

REFERENCES

1. S. H. Brecht, D. A. Hitchcock, and W. Horton, Jr., Phys. Fluids 21, 447, (1978).
2. D. A. Hitchcock, S. H. Brecht, and W. Horton, Jr., Phys. Fluids 20, 1551, (1977).
3. E. S. Weibel, Phys. Fluids 13, 3003, (1970).
4. H. L. Berk, W. Horton, Jr., M. N. Rosenbluth, and P. H. Rutherford, Nuclear Fusion 15, 1012 (1976).
5. L. P. Mai and W. Horton, Jr., Phys. Fluids 18, 356, (1975).
6. M. Yamada, S. Seiler, H. W. Hendel and H. Ikeji, Phys. Fluids 20, 450, (1977).
7. H. Böhmer, J. P. Hauch and N. Ryan, Phys. Fluids 19, 450 (1976).
8. H. Böhmer, Phys. Fluids 19, 1371, (1976).
9. P. Michelen, H. L. Aecseli, J. J. Rasmussen, N. Sato, Phys. Fluids 19, 453, (1976).
10. R. Dei-Cas in "Third Symposium on Plasma Heating in Toroidal Devices" (Proceeding of the International School of Plasma Physics, Villa Monastero, Italy, 1976). Editor E. Sindoni, Editrice Compositori, Bologna (1976) pp. 781-789.
11. S. H. Brecht, Univ. of Texas, FRCR No. 163 (1977).
12. N. A. Krall and A. W. Trivelpiece, "Principles of Plasma Physics (McGraw-Hill, New York, 1973).
13. G. T. Davidson, JGR 78, 7569 (1973).
14. G. T. Davidson, JGR 80, 3172 (1975).

Table I
Input Parameters for Ion Bernstein Mode

Electron Temperature	0.07 eV
Ion Temperature	0.07 eV
Magnetic Field	0.3 G
Electron Density	$1.0 \times 10^4/\text{cm}^3$
Mass of Beam Particle	$27 m_p$
Beam Density	$0.2 n_e$
Mass of Background Ions	$16 m_p$
Initial Beam Velocity	$1.18 \times 10^8 \text{ cm/sec}$ or 200 keV

Table II

Results for the second Harmonic Excitation

l	$= 2$
Growth Rate γ ,	$\leq 2\omega_{ci}$
$\omega - l\omega_{ci}$	$\sim 0.01\omega_{ci} \rightarrow 0.01 \text{ ci}$
ω_{ci}	$\sim 1.80 \times 10^2/\text{sec}$
ω_{cb}	$\sim 1.06 \times 10^2/\text{sec}$
ω	$\sim 3.56 \times 10^2/\text{sec}$
k_{\perp}	$\sim 1.E-6/\text{cm}$
$k_{ }$	$\sim 2.E-6/\text{cm}$
v_{thi}	$\sim 9.15 \times 10^4 \text{ cm/sec}$
v_{gi}	$\sim 2.97 \times 10^2 \text{ cm/sec}$
$v_{g }$	$\sim 1 \times 10^7 \text{ cm/sec}$

Table III

Parameter for Quasilinear Estimate

$\Delta v_{b\perp}$	\sim	$v_{b\perp} \sim \ell \omega_{cb} / k_{\perp}$
$\Delta v_{b }$	\sim	$0.2 v_{b }$
$v_{b\perp}$	\sim	$0.1 v_{b }$
$\Delta k_{ }$	\sim	$3 \times 10^{-7} / \text{cm}$
Δk_{\perp}	\sim	$3 \times 10^{-6} / \text{cm}$
$J_{\ell}(b) e^{-b}$	\sim	$(b/2)^{\ell} e^{-b} \sim (b/2)^{\ell}$
$J_{\ell}(k_{\perp} v_{\perp} / \omega_{cb})$	\sim	$(1/2 k_{\perp} v_{\perp} / \omega_{cb})^{\ell}$
$\Delta^3 k$	\sim	$\Delta k_{ } \Delta k_{\perp}^2$
c_s	\sim	$6.47 \times 10^5 \text{ cm/sec}$
v_b	\sim	$1.18 \times 10^8 \text{ cm/sec}$




Figure 1

The debris distribution functions as used in the linear calculation. The velocity of injection is $1.18 \times 10^8 \text{ cm/sec}$. The velocities on the plot are in dimensionless units. The normalization velocity is $5.59 \times 10^6 \text{ cm/sec}$.




Figure 2

The time evolution of the perpendicular velocity of the streaming debris. The initial $v_{b\perp}$ is assumed 0.1 of $v_{b\parallel}$. The parallel correlation time, τ_c , and τ_s , the estimated time for parallel trapping effects are shown for each mode. In both $\ell = 2$ and $\ell = 3$ case it was assumed that $\gamma \sim \ell \omega_{ci}$.

DISTRIBUTION LIST

DEPARTMENT OF DEFENSE

ASSISTANT SECRETARY OF DEFENSE
COMM, CMD, CONT & INTELL
WASHINGTON, D.C. 20301
01CY ATTN J. BABCOCK
01CY ATTN M. EPSTEIN

ASSISTANT TO THE SECRETARY OF DEFENSE
ATOMIC ENERGY
WASHINGTON, D.C. 20301
01CY ATTN EXECUTIVE ASSISTANT

DIRECTOR
COMMAND CONTROL TECHNICAL CENTER
PENTAGON RM BE 685
WASHINGTON, D.C. 20301
01CY ATTN C-650
01CY ATTN C-312 R. MASON

DIRECTOR
DEFENSE ADVANCED RSCH PROJ AGENCY
ARCHITECT BUILDING
1400 WILSON BLVD.
ARLINGTON, VA. 22209
01CY ATTN NUCLEAR MONITORING RESEARCH
01CY ATTN STRATEGIC TECH OFFICE

DEFENSE COMMUNICATION ENGINEER CENTER
1860 WIEHLE AVENUE
RESTON, VA. 22090
01CY ATTN CODE R820
01CY ATTN CODE R410 JAMES W. MCLEAN
01CY ATTN CODE R720 J. WORTHINGTON

DIRECTOR
DEFENSE COMMUNICATIONS AGENCY
WASHINGTON, D.C. 20305
(ADR CNWDI: ATTN CODE 240 FOR)
01CY ATTN CODE 101B

DEFENSE DOCUMENTATION CENTER
CAMERON STATION
ALEXANDRIA, VA. 22314
(12 COPIES IF OPEN PUBLICATION, OTHERWISE 2 COPIES)
12CY ATTN TC

DIRECTOR
DEFENSE INTELLIGENCE AGENCY
WASHINGTON, D.C. 20301
01CY ATTN DT-1B
01CY ATTN DB-4C E. O'FARRELL
01CY ATTN DIAAP A. WISE
01CY ATTN DIAST-5
01CY ATTN DT-1BZ R. MORTON
01CY ATTN HQ-TR J. STEWART
01CY ATTN W. WITTIG DC-7D

DIRECTOR
DEFENSE NUCLEAR AGENCY
WASHINGTON, D.C. 20305
01CY ATTN STVL
04CY ATTN TITL
01CY ATTN DOST
03CY ATTN RAAE

COMMANDER
FIELD COMMAND
DEFENSE NUCLEAR AGENCY
KIRTLAND AFB, NM 87115
01CY ATTN FCPR

DIRECTOR
INTERSERVICE NUCLEAR WEAPONS SCHOOL
KIRTLAND AFB, NM 87115
01CY ATTN DOCUMENT CONTROL

JOINT CHIEFS OF STAFF
WASHINGTON, D.C. 20301
01CY ATTN J-3 WMMCCS EVALUATION OFFICE

DIRECTOR
JOINT STRAT TGT PLANNING STAFF
OFFUTT AFB
OMAHA, NB 68113
01CY ATTN ULTW-2
01CY ATTN JPST G. GOETZ

CHIEF
LIVERMORE DIVISION FLD COMMAND DNA
DEPARTMENT OF DEFENSE
LAWRENCE LIVERMORE LABORATORY
P. O. BOX 808
LIVERMORE, CA 94550
01CY ATTN FCPLR

DIRECTOR
NATIONAL SECURITY AGENCY
DEPARTMENT OF DEFENSE
FT. GEORGE G. MEADE, MD 20755
01CY ATTN JOHN SKILLMAN R52
01CY ATTN FRANK LEONARD
01CY ATTN W14 PAT CLARK
01CY ATTN OLIVER M. BARTLETT W32
01CY ATTN R5

COMMANDANT
NATO SCHOOL (SHAPE)
APO NEW YORK 09172
01CY ATTN U.S. DOCUMENTS OFFICER

UNDER SECY OF DEF FOR RSCH & ENGRG
DEPARTMENT OF DEFENSE
WASHINGTON, D.C. 20301
01CY ATTN STRATEGIC & SPACE SYSTEMS (OS)

WMMCCS SYSTEM ENGINEERING ORG
WASHINGTON, D.C. 20305
01CY ATTN R. CRAWFORD

COMMANDER/DIRECTOR
ATMOSPHERIC SCIENCES LABORATORY
U.S. ARMY ELECTRONICS COMMAND
WHITE SANDS MISSILE RANGE, NM 88002
01CY ATTN DELAS-EO F. NILES

DIRECTOR
BMD ADVANCED TECH CTR
HUNTSVILLE OFFICE
P. O. BOX 1500
HUNTSVILLE, AL 35807
01CY ATTN ATC-T MELVIN T. CAPPS
01CY ATTN ATC-O W. DAVIES
01CY ATTN ATC-R DON RUSS

PROGRAM MANAGER
BMD PROGRAM OFFICE
5001 EISENHOWER AVENUE
ALEXANDRIA, VA 22333
OICY ATTN DACS-BMT J. SHEA

CHIEF C-E SERVICES DIVISION
U.S. ARMY COMMUNICATIONS CMD
PENTAGON RM 18269
WASHINGTON, D.C. 20310
OICY ATTN C-E-SERVICES DIVISION

COMMANDER
FRADCOM TECHNICAL SUPPORT ACTIVITY
DEPARTMENT OF THE ARMY
FORT MONMOUTH, N.J. 07703
OICY ATTN DRSEL-NL-RD M. BENNET
OICY ATTN DRSEL-PL-ENV M. BOMKE
OICY ATTN J. E. QUIGLEY

COMMANDER
HARRY DIAMOND LABORATORIES
DEPARTMENT OF THE ARMY
2800 POWDER MILL ROAD
ADELPHI, MD 20783
(CNWDI-INNER ENVELOPE: ATTN: DELHD-RBM)
OICY ATTN DELHD-TI M. WEINER
OICY ATTN DELHD-RB R. WILLIAMS
OICY ATTN DELHD-NP F. WIMENITZ
OICY ATTN DELHD-NP C. MOAZED

COMMANDER
U.S. ARMY COMM-ELEC ENGRG INSTAL AGY
FT. HUACHUCA, AZ 85613
OICY ATTN CCC-EMEO GEORGE LANE

COMMANDER
U.S. ARMY FOREIGN SCIENCE & TECH CTR
220 7TH STREET, NE
CHARLOTTESVILLE, VA 22901
OICY ATTN DRXST-SD
OICY ATTN R. JONES

COMMANDER
U.S. ARMY MATERIEL DEV & READINESS CMD
5001 EISENHOWER AVENUE
ALEXANDRIA, VA 22333
OICY ATTN DRCLDC J. A. BENDER

COMMANDER
U.S. ARMY NUCLEAR AND CHEMICAL AGENCY
7500 BACKLICK ROAD
BLDG 2073
SPRINGFIELD, VA 22150
OICY ATTN LIBRARY

DIRECTOR
U.S. ARMY BALLISTIC RESEARCH LABS
ABERDEEN PROVING GROUND, MD 21005
OICY ATTN TECH LIB EDWARD BAICY

COMMANDER
U.S. ARMY SATCOM AGENCY
FT. MONMOUTH, NJ 07703
OICY ATTN DOCUMENT CONTROL

COMMANDER
U.S. ARMY MISSILE INTELLIGENCE AGENCY
REDSTONE ARSENAL, AL 35809
OICY ATTN JIM GAMBLE

DIRECTOR
U.S. ARMY TRADOC SYSTEMS ANALYSIS ACTIVITY
WHITE SANDS MISSILE RANGE, NM 88002
OICY ATTN ATAA-SA
OICY ATTN TCC/F. PAYAN JR.
OICY ATTN ATAA-TAC LTC J. HESSE

COMMANDER
NAVAL ELECTRONIC SYSTEMS COMMAND
WASHINGTON, D.C. 20360
OICY ATTN NAVALEX 034 T. HUGHES
OICY ATTN PME 117
OICY ATTN PME 117-T
OICY ATTN CODE 5011

COMMANDING OFFICER
NAVAL INTELLIGENCE SUPPORT CTR
4301 SUITLAND ROAD, BLDG. 5
WASHINGTON, D.C. 20390
OICY ATTN MR. DUBBIN STIC 12
OICY ATTN NISC-50
OICY ATTN CODE 5404 J. GALET

COMMANDER
NAVAL OCEAN SYSTEMS CENTER
SAN DIEGO, CA 92152
OICY ATTN CODE 532 W. MOLER
OICY ATTN CODE 0230 C. BAGGETT
OICY ATTN CODE 81 R. EASTMAN

DIRECTOR
NAVAL RESEARCH LABORATORY
WASHINGTON, D.C. 20375
OICY ATTN CODE 4700 TIMOTHY P. COFFEY (25 CYS
IF UNCLASS, 1 CY IF CLASS)
OICY ATTN CODE 4701 JACK D. BROWN
OICY ATTN CODE 4780 BRANCH HEAD (150 CYS
IF UNCLASS, 1 CY IF CLASS)
OICY ATTN CODE 7500 HQ COMM DIR BRUCE WALD
OICY ATTN CODE 7550 J. DAVIS
OICY ATTN CODE 7580
OICY ATTN CODE 7551
OICY ATTN CODE 7555
OICY ATTN CODE 4730 E. MCLEAN
OICY ATTN CODE 4127 C. JOHNSON

COMMANDER
NAVAL SEA SYSTEMS COMMAND
WASHINGTON, D.C. 20362
OICY ATTN CAPT R. PITKIN

COMMANDER
NAVAL SPACE SURVEILLANCE SYSTEM
DAHLGREN, VA 22448
OICY ATTN CAPT J. H. BURTON

OFFICER-IN-CHARGE
NAVAL SURFACE WEAPONS CENTER
WHITE OAK, SILVER SPRING, MD 20910
OICY ATTN CODE F31

DIRECTOR
STRATEGIC SYSTEMS PROJECT OFFICE
DEPARTMENT OF THE NAVY
WASHINGTON, D.C. 20376
OICY ATTN NSP-2141
OICY ATTN NSSP-2722 FRED WIMBERLY

NAVAL SPACE SYSTEM ACTIVITY
P. O. BOX 92960
WORLDWAY POSTAL CENTER
LOS ANGELES, CALIF. 90009
OICY ATTN A. B. HAZZARD

COMMANDER
NAVAL SURFACE WEAPONS CENTER
DAHLGREN LABORATORY
DAHLGREN, VA 22448
OICY ATTN CODE DF-14 R. BUTLER

COMMANDING OFFICER
NAVY SPACE SYSTEMS ACTIVITY
P.O. BOX 92960
WORLDWAY POSTAL CENTER
LOS ANGELES, CA. 90009
OICY ATTN CODE 52

OFFICE OF NAVAL RESEARCH
ARLINGTON, VA 22217
OICY ATTN CODE 465
OICY ATTN CODE 461
OICY ATTN CODE 402
OICY ATTN CODE 420
OICY ATTN CODE 421

COMMANDER
AEROSPACE DEFENSE COMMAND/DC
DEPARTMENT OF THE AIR FORCE
ENT AFB, CO 80912
OICY ATTN DC MR. LONG

COMMANDER
AEROSPACE DEFENSE COMMAND/XPD
DEPARTMENT OF THE AIR FORCE
ENT AFB, CO 80912
OICY ATTN XPDQ
OICY ATTN XP

AIR FORCE GEOPHYSICS LABORATORY
HANSOM AFB, MA 01731
OICY ATTN OPR HAROLD GARDNER
OICY ATTN OPR-1 JAMES C. ULWICK
OICY ATTN LKB KENNETH S. W. CHAMPION
OICY ATTN OPR ALVA T. STAIR
OICY ATTN PHP JULES AARONS
OICY ATTN PHD JURGEN BUCHAU
OICY ATTN PHD JOHN P. MULLEN

AF WEAPONS LABORATORY
KIRTLAND AFB, NM 87117
OICY ATTN SUL
OICY ATTN CA ARTHUR H. GUENTHER
OICY ATTN DYC CAPT J. BARRY
OICY ATTN DYC JOHN M. KAHN
OICY ATTN DYT CAPT MARK A. FRY
OICY ATTN DES MAJ GARY GANDONG
OICY ATTN DYC J. JANNI

AFTAC
PATRICK AFB, FL 32925
OICY ATTN TF/MAJ WILEY
OICY ATTN TN

AIR FORCE AVIONICS LABORATORY
WRIGHT-PATTERSON AFB, OH 45433
OICY ATTN AAD WADE HUNT
OICY ATTN AAD ALLEN JOHNSON

DEPUTY CHIEF OF STAFF
RESEARCH, DEVELOPMENT, & ACQ
DEPARTMENT OF THE AIR FORCE
WASHINGTON, D.C. 20330
OICY ATTN AFRDQ

HEADQUARTERS
ELECTRONIC SYSTEMS DIVISION/XR
DEPARTMENT OF THE AIR FORCE
HANSOM AFB, MA 01731
OICY ATTN XR J. DEAS

HEADQUARTERS
ELECTRONIC SYSTEMS DIVISION/YSEA
DEPARTMENT OF THE AIR FORCE
HANSOM AFB, MA 01731
OICY ATTN YSEA

HEADQUARTERS
ELECTRONIC SYSTEMS DIVISION/DC
DEPARTMENT OF THE AIR FORCE
HANSOM AFB, MA 01731
OICY ATTN DCKC MAJ J.C. CLARK

COMMANDER
FOREIGN TECHNOLOGY DIVISION, AFSC
WRIGHT-PATTERSON AFB, OH 45433
OICY ATTN NICD LIBRARY
OICY ATTN ETD P. BALLARD

COMMANDER
ROME AIR DEVELOPMENT CENTER, AFSC
GRIFFISS AFB, NY 13441
OICY ATTN DOC LIBRARY/TSLD
OICY ATTN OCSE V. COYNE

SAMSO/SZ
POST OFFICE BOX 92960
WORLDWAY POSTAL CENTER
LOS ANGELES, CA 90009
(SPACE DEFENSE SYSTEMS)
OICY ATTN SZJ

STRATEGIC AIR COMMAND/XPFS
OFFUTT AFB, NE 68113
OICY ATTN XPFS MAJ B. STEPHAN
OICY ATTN ADMATE MAJ BRUCE BAUER
OICY ATTN NRT
OICY ATTN DOK CHIEF SCIENTIST

SAMSO/YA
P. O. BOX 92960
WORLDWAY POSTAL CENTER
LOS ANGELES, CA 90009
OICY ATTN YAT CAPT L. BLACKWELDER

SAMSO/SK
P. O. BOX 92960
WORLDWAY POSTAL CENTER
LOS ANGELES, CA 90009
OICY ATTN SKA (SPACE COMM SYSTEMS) M. CLAVIN

SAMSO/MN
NORTON AFB, CA 92409
(MINUTEMAN)
OICY ATTN MNML LTC KENNEDY

COMMANDER
ROME AIR DEVELOPMENT CENTER, AFSC
HANSOM AFB, MA 01731
OICY ATTN EEP A. LORENTZEN

DEPARTMENT OF ENERGY
ALBUQUERQUE OPERATIONS OFFICE
P. O. BOX 5400
ALBUQUERQUE, NM 87115
OICY ATTN DOC CON FOR D. SHERWOOD

DEPARTMENT OF ENERGY
LIBRARY ROOM G-042
WASHINGTON, D.C. 20545
OICY ATTN DOC CON FOR A. LABOWITZ

EG&G, INC.
LOS ALAMOS DIVISION
P. O. BOX 809
LOS ALAMOS, NM 85544
01CY ATTN DOC CON FOR J. BREEDLOVE

UNIVERSITY OF CALIFORNIA
LAWRENCE LIVERMORE LABORATORY
P. O. BOX 808
LIVERMORE, CA 94550
01CY ATTN DOC CON FOR TECH INFO DEPT
01CY ATTN DOC CON FOR L-389 R. OTT
01CY ATTN DOC CON FOR L-31 R. HAGER
01CY ATTN DOC CON FOR L-46 F. SEWARD

LOS ALAMOS SCIENTIFIC LABORATORY
P. O. BOX 1663
LOS ALAMOS, NM 87545
01CY ATTN DOC CON FOR J. WOLCOTT
01CY ATTN DOC CON FOR R. F. TASCHER
01CY ATTN DOC CON FOR E. JONES
01CY ATTN DOC CON FOR J. MALIK
01CY ATTN DOC CON FOR R. JEFFRIES
01CY ATTN DOC CON FOR J. ZINN
01CY ATTN DOC CON FOR P. KEATON
01CY ATTN DOC CON FOR D. WESTERVELT

SANDIA LABORATORIES
P. O. BOX 5800
ALBUQUERQUE, NM 87115
01CY ATTN DOC CON FOR J. MARTIN
01CY ATTN DOC CON FOR W. BROWN
01CY ATTN DOC CON FOR A. THORNBROUGH
01CY ATTN DOC CON FOR T. WRIGHT
01CY ATTN DOC CON FOR D. DAHLGREN
01CY ATTN DOC CON FOR 3141
01CY ATTN DOC CON FOR SPACE PROJECT DIV

SANDIA LABORATORIES
LIVERMORE LABORATORY
P. O. BOX 969
LIVERMORE, CA 94550
01CY ATTN DOC CON FOR B. MURPHEY
01CY ATTN DOC CON FOR T. COOK

OFFICE OF MILITARY APPLICATION
DEPARTMENT OF ENERGY
WASHINGTON, D.C. 20545
01CY ATTN DOC CON FOR D. GALE

OTHER GOVERNMENT

CENTRAL INTELLIGENCE AGENCY
ATTN RD/51, RM 5G48, HQ BLDG
WASHINGTON, D.C. 20505
01CY ATTN OSI/PSID RM 5F 19

DEPARTMENT OF COMMERCE
NATIONAL BUREAU OF STANDARDS
WASHINGTON, D.C. 20234
(CALL CORRES: ATTN SEC OFFICER FOR)
01CY ATTN R. MOORE

INSTITUTE FOR TELECOM SCIENCES
NATIONAL TELECOMMUNICATIONS & INFO ADMIN
BOULDER, CO 80303
01CY ATTN A. JEAN (UNCLASS ONLY)
01CY ATTN W. UTLAUT
01CY ATTN D. CROMBIE
01CY ATTN L. BERRY

NATIONAL OCEANIC & ATMOSPHERIC ADMIN
ENVIRONMENTAL RESEARCH LABORATORIES
DEPARTMENT OF COMMERCE
BOULDER, CO 80302
01CY ATTN R. GRUBB
01CY ATTN AERONOMY LAB G. REID

DEPARTMENT OF DEFENSE CONTRACTORS

AEROSPACE CORPORATION
P. O. BOX 92957
LOS ANGELES, CA 90009
01CY ATTN I. GARFUNKEL
01CY ATTN T. SALMI
01CY ATTN V. JOSEPHSON
01CY ATTN S. BOWER
01CY ATTN N. STOCKWELL
01CY ATTN D. OLSEN
01CY ATTN J. CARTER
01CY ATTN F. MORSE
01CY ATTN SMFA FOR PMW

ANALYTICAL SYSTEMS ENGINEERING CORP
5 OLD CONCORD ROAD
BURLINGTON, MA 01803
01CY ATTN RADIO SCIENCES

BERKELEY RESEARCH ASSOCIATES, INC.
P. O. BOX 983
BERKELEY, CA 94701
01CY ATTN J. WORKMAN

BOEING COMPANY, THE
P. O. BOX 3707
SEATTLE, WA 98124
01CY ATTN G. KEISTER
01CY ATTN D. MURRAY
01CY ATTN G. HALL
01CY ATTN J. KENNEY

CALIFORNIA AT SAN DIEGO, UNIV OF
IPAPS, B-019
LA JOLLA, CA 92093
01CY ATTN HENRY G. BOOKER

BROWN ENGINEERING COMPANY, INC.
CUMMINGS RESEARCH PARK
HUNTSVILLE, AL 35807
01CY ATTN ROMEO A. DELIBERIS

CHARLES STARK DRAPER LABORATORY, INC.
555 TECHNOLOGY SQUARE
CAMBRIDGE, MA 02139
01CY ATTN D. B. COX
01CY ATTN J. P. GILMORE

COMPUTER SCIENCES CORPORATION
6565 ARLINGTON BLVD
FALLS CHURCH, VA 22046
01CY ATTN H. BLANK
01CY ATTN JOHN SPOOR
01CY ATTN C. NAIL

COMSAT LABORATORIES
LINTHICUM ROAD
CLARKSBURG, MD 20734
01CY ATTN G. HYDE

CORNELL UNIVERSITY
DEPARTMENT OF ELECTRICAL ENGINEERING
ITHACA, NY 14850
01CY ATTN D. T. FARLEY JR

ELECTROSPACE SYSTEMS, INC.
BOX 1359
RICHARDSON, TX 75080
OICY ATTN H. LOGSTON
OICY ATTN SECURITY (PAUL PHILLIPS)

ESL INC.
495 JAVA DRIVE
SUNNYVALE, CA 94086
OICY ATTN J. ROBERTS
OICY ATTN JAMES MARSHALL
OICY ATTN C. W. PRETTIE

FORD AEROSPACE & COMMUNICATIONS CORP
3939 FABIAN WAY
PALO ALTO, CA 94303
OICY ATTN J. T. MATTINGLEY

GENERAL ELECTRIC COMPANY
SPACE DIVISION
VALLEY FORGE SPACE CENTER
GODDARD BLVD KING OF PRUSSIA
P. O. BOX 8555
PHILADELPHIA, PA 19101
OICY ATTN M. H. BORTNER SPACE SCI LAB

GENERAL ELECTRIC COMPANY
P. O. BOX 1122
SYRACUSE, NY 13201
OICY ATTN F. REIBERT

GENERAL ELECTRIC COMPANY
TEMPO-CENTER FOR ADVANCED STUDIES
816 STATE STREET (P.O. DRAWER QQ)
SANTA BARBARA, CA 93102
OICY ATTN DASAC
OICY ATTN DON CHANDLER
OICY ATTN TOM BARRETT
OICY ATTN TIM STEPHANS
OICY ATTN WARREN S. KNAPP
OICY ATTN WILLIAM McNAMARA
OICY ATTN B. GAMBILL
OICY ATTN MACK STANTON

GENERAL ELECTRIC TECH SERVICES CO., INC.
HMES
COURT STREET
SYRACUSE, NY 13201
OICY ATTN G. MILLMAN

GENERAL RESEARCH CORPORATION
SANTA BARBARA DIVISION
P. O. BOX 6770
SANTA BARBARA, CA 93111
OICY ATTN JOHN ISE JR
OICY ATTN JOEL GARBARINO

GEOPHYSICAL INSTITUTE
UNIVERSITY OF ALASKA
FAIRBANKS, AK 99701
(ALL CLASS ATTN: SECURITY OFFICER)
OICY ATTN T. N. DAVIS (UNCL ONLY)
OICY ATTN NEAL BROWN (UNCL ONLY)
OICY ATTN TECHNICAL LIBRARY

GTE SYLVANIA, INC.
ELECTRONICS SYSTEMS GRP-EASTERN DIV
77 A STREET
NEEDHAM, MA 02194
OICY ATTN MARSHAL CROSS

ILLINOIS, UNIVERSITY OF
DEPARTMENT OF ELECTRICAL ENGINEERING
URBANA, IL 61803
OICY ATTN K. YEH

ILLINOIS, UNIVERSITY OF
107 COBLE HALL
801 S. WRIGHT STREET
URBANA, IL 60680
(ALL CORRES ATTN SECURITY SUPERVISOR FOR)
OICY ATTN K. YEH

INSTITUTE FOR DEFENSE ANALYSES
400 ARMY-NAVY DRIVE
ARLINGTON, VA 22202
OICY ATTN J. M. AEIN
OICY ATTN ERNEST BAUER
OICY ATTN HANS WOLFHARD
OICY ATTN JOEL BENGSTON

HSS, INC.
2 ALFRED CIRCLE
BEDFORD, MA 01730
OICY ATTN DONALD HANSEN

INTL TEL & TELEGRAPH CORPORATION
500 WASHINGTON AVENUE
NUTLEY, NJ 07110
OICY ATTN TECHNICAL LIBRARY

JAYCOR
1401 CAMINO DEL MAR
DEL MAR, CA 92014

JOHNS HOPKINS UNIVERSITY
APPLIED PHYSICS LABORATORY
JOHNS HOPKINS ROAD
LAUREL, MD 20810
OICY ATTN DOCUMENT LIBRARIAN
OICY ATTN THOMAS POTEHRA
OICY ATTN JOHN DASSOULAS

LOCKHEED MISSILES & SPACE CO INC
P. O. BOX 504
SUNNYVALE, CA 94088
OICY ATTN DEPT 60-12
OICY ATTN D. R. CHURCHILL

LOCKHEED MISSILES AND SPACE CO INC
3251 MANOVER STREET
PALO ALTO, CA 94304
OICY ATTN MARTIN WALT DEPT 52-10
OICY ATTN RICHARD G. JOHNSON DEPT 52-12
OICY ATTN W. L. IMHOFF DEPT 52-12

KAMAN SCIENCES CORP
P. O. BOX 7463
COLORADO SPRINGS, CO 80933
OICY ATTN T. MEAGHER

LINKABIT CORP
10453 ROSELLE
SAN DIEGO, CA 92121
OICY ATTN IRWIN JACOBS

M.I.T. LINCOLN LABORATORY
P. O. BOX 73
LEXINGTON, MA 02173
OICY ATTN DAVID M. TOWLE
OICY ATTN P. WALDRON
OICY ATTN L. LOUGHLIN
OICY ATTN D. CLARK

MARTIN MARIETTA CORP
ORLANDO DIVISION
P. O. BOX 5837
ORLANDO, FL 32805
OICY ATTN R. HEFFNER

MCDONNELL DOUGLAS CORPORATION
5301 BOLSA AVENUE
HUNTINGTON BEACH, CA 92647
OICY ATTN N. HARRIS
OICY ATTN J. MOULE
OICY ATTN GEORGE MROZ
OICY ATTN W. OLSON
OICY ATTN R. W. HALPRIN
OICY ATTN TECHNICAL LIBRARY SERVICES

MISSION RESEARCH CORPORATION
735 STATE STREET
SANTA BARBARA, CA 93101
OICY ATTN P. FISCHER
OICY ATTN W. F. CREVIER
OICY ATTN STEVEN L. GUTSCHE
OICY ATTN D. SAPPENFELD
OICY ATTN R. BOGUSCH
OICY ATTN R. MENDRICK
OICY ATTN RALPH KILB
OICY ATTN DAVE SOWLE
OICY ATTN F. FAJEN
OICY ATTN M. SCHEIBE
OICY ATTN CONRAD L. LONGMIRE
OICY ATTN WARREN A. SCHLUTER

MITRE CORPORATION, THE
P. O. BOX 208
BEDFORD, MA 01730
OICY ATTN JOHN MORGANSTERN
OICY ATTN G. HARDING
OICY ATTN C. E. CALLAHAN

MITRE CORP
WESTGATE RESEARCH PARK
1820 DOLLY MADISON BLVD
MCLEAN, VA 22101
OICY ATTN W. HALL
OICY ATTN W. FOSTER

PACIFIC-SIERRA RESEARCH CORP
1456 CLOVERFIELD BLVD.
SANTA MONICA, CA 90404
OICY ATTN E. C. FIELD JR

PENNSYLVANIA STATE UNIVERSITY
IONOSPHERE RESEARCH LAB
318 ELECTRICAL ENGINEERING EAST
UNIVERSITY PARK, PA 16802
(NO CLASSIFIED TO THIS ADDRESS)
OICY ATTN IONOSPHERIC RESEARCH LAB

PHOTOMETRICS, INC.
442 MARRETT ROAD
LEXINGTON, MA 02173
OICY ATTN IRVING L. KOFSKY

PHYSICAL DYNAMICS INC.
P. O. BOX 3027
BELLEVUE, WA 98009
OICY ATTN E. J. FREMOW

PHYSICAL DYNAMICS INC.
P. O. BOX 1069
BERKELEY, CA 94701
OICY ATTN A. THOMPSON

R & D ASSOCIATES
P. O. BOX 9695
MARINA DEL REY, CA 90291
OICY ATTN FORREST GILMORE
OICY ATTN BRYAN GABBARD
OICY ATTN WILLIAM B. WRIGHT JR
OICY ATTN ROBERT F. LELEVIER
OICY ATTN WILLIAM J. KARZAS
OICY ATTN H. ORY
OICY ATTN C. MACDONALD
OICY ATTN R. TURCO

RAND CORPORATION, THE
1700 MAIN STREET
SANTA MONICA, CA 90406
OICY ATTN CULLEN CRAIN
OICY ATTN ED BEDROZIAN

RIVERSIDE RESEARCH INSTITUTE
80 WEST END AVENUE
NEW YORK, NY 10023
OICY ATTN VINCE TRAPANI

SCIENCE APPLICATIONS, INC.
P. O. BOX 2351
LA JOLLA, CA 92038
OICY ATTN LEWIS M. LINSON
OICY ATTN DANIEL A. HAMLIN
OICY ATTN D. SACHS
OICY ATTN E. A. STRAKER
OICY ATTN CURTIS A. SMITH
OICY ATTN JACK MCDUGALL

RAYTHEON CO.
528 BOSTON POST ROAD
SUDBURY, MA 01776
OICY ATTN BARBARA ADAMS

SCIENCE APPLICATIONS, INC.
HUNTSVILLE DIVISION
2109 W. CLINTON AVENUE
SUITE 700
HUNTSVILLE, AL 35805
OICY ATTN DALE H. DIVIS

SCIENCE APPLICATIONS, INCORPORATED
8400 WESTPARK DRIVE
MCLEAN, VA 22101
OICY ATTN J. COCKAYNE

SCIENCE APPLICATIONS, INC.
80 MISSION DRIVE
PLEASANTON, CA 94566
OICY ATTN SZ

SRI INTERNATIONAL
333 RAVENSWOOD AVENUE
MENLO PARK, CA 94025
OICY ATTN DONALD NEILSON
OICY ATTN ALAN BURNS
OICY ATTN G. SMITH
OICY ATTN L. L. COBB
OICY ATTN DAVID A. JOHNSON
OICY ATTN WALTER G. CHESNUT
OICY ATTN CHARLES L. RIND
OICY ATTN WALTER JAYE
OICY ATTN M. BARON
OICY ATTN RAY L. LEADABRAND
OICY ATTN G. CARPENTER
OICY ATTN G. PRICE
OICY ATTN J. PETERSON
OICY ATTN R. HAKE, JR.
OICY ATTN V. GONZALES
OICY ATTN D. MCDANIEL

TECHNOLOGY INTERNATIONAL CORP
75 WIGGINS AVENUE
BEDFORD, MA 01730
OICY ATTN W. P. BOQUIST

TRW DEFENSE & SPACE SYS GROUP
ONE SPACE PARK
REDONDO BEACH, CA 90278
OICY ATTN R. K. PLEBUCH
OICY ATTN S. ALTSCHULER
OICY ATTN D. DEE

VISIODYNE, INC.
19 THIRD AVENUE
NORTH WEST INDUSTRIAL PARK
BURLINGTON, MA 01803
OICY ATTN CHARLES HUMPHREY
OICY ATTN J. W. CARPENTER